

LMA MSW Solution in the Minimal $SU(3)_L \times U(1)_N$ Gauge Model [★]

Teruyuki Kitabayashi^a, Masaki Yasuè^{b,c}

^a Accelerator Engineering Center,
Mitsubishi Electric System & Service Engineering Co.Ltd.,
2-8-8 Umezono, Tsukuba, Ibaraki 305-0045, Japan.

^b Department of Natural Science,
School of Marine Science and Technology, Tokai University,
3-20-1 Orido, Shimizu, Shizuoka 424-8610, Japan

^c Department of Physics, Tokai University,
1117 Kitakaname, Hiratsuka, Kanagawa 259-1291, Japan.

Abstract

The minimal $SU(3)_L \times U(1)_N$ gauge model for (ν, ℓ, ℓ^C) equipped with the (approximate) $L_e - L_\mu - L_\tau$ symmetry and a discrete Z_4 symmetry is found to provide radiative neutrino masses compatible with the LMA MSW solution.

The neutrino oscillation phenomena imply the existence of massive neutrinos [1]. Non-zero Majorana neutrino masses are known to arise in models with lepton number violating interactions and $SU(3)_L \times U(1)_N$ models naturally have such interactions [2], which are linked to generate charged lepton masses. With this in mind, we have constructed the simplest $SU(3)_L \times U(1)_N$ model with for (ν, ℓ, ℓ^C) giving rise to the currently most favorable LMA solution [3].

We use the following three guidelines to construct our model. (1) the model accounts for $\Delta m_{atm}^2 \gg \Delta m_\odot^2$ [4], which will be realized by bimaximal neutrino mixings based on the (approximate) $L' \equiv L_e - L_\mu - L_\tau$ symmetry [5]; (2) the model respects the LMA solution, $\Delta m_{atm}^2 / \Delta m_\odot^2 = 10^{-2} - 10^{-1}$ [4]; and (3) the model provides neutrino masses as radiative effects [6].

The particle content is given by, for $(U(1)_N, L')$: (a) triplet leptons, $\psi^1 = (\nu^1, \ell^1, \ell^{c1})_L^T : (0, 1)$, $\psi^{i=2,3} = (\nu^i, \ell^i, \ell^{ci})_L^T : (0, -1)$; (b) triplet scalars, $\eta = (\eta^0, \eta^-, \eta^+)^T : (0, 0)$, $\rho = (\rho^+, \rho^0, \rho^{++})^T : (1, 0)$, $\chi = (\chi^-, \chi^{--}, \chi^0)^T : (-1, 0)$

[★] Talk given by T.K. at the International Workshop NuFACT'01, Tsukuba, Japan (May 2001).




Fig. 1. One-loop diagram for δm_{11}^{rad} , where $m_{\ell i} = f_{ii} v_{\ell}^{(+)}$.

with $\langle \eta^0, \rho^0, \chi^0 \rangle (\equiv v_{\eta, \rho, \chi})$ and (c) antisextet scalars, $S^{(0)} : (1, 0)$ for $\psi^1 \psi^{2,3}$, $S^{(+)} : (1, 2)$ for $\psi^{2,3} \psi^{2,3}$. If there is only one antisextet, because the antisextet yields both mass terms for ℓ and ν , we will encounter the similar flavor mass-structure for ν and ℓ , which, however, contradicts with the phenomenology. To avoid this 'similarity', we introduce two antisextets of $S^{(0,+)\alpha\beta}$ ($\alpha, \beta=1,2,3$) and study their vacuum alignment to yield, for neutrinos, $\langle S^{(0)11} \rangle (\equiv v_{\nu}^{(0)})$ but without $\langle S^{(+11)} \rangle (\equiv v_{\nu}^{(+)})$ and, for charged leptons, $\langle S^{(0,+23)} \rangle (\equiv v_{\ell}^{(0,+)})$ [3].

To realize the vacuum alignment, we use the L' symmetry with a Z_4 symmetry, where $Z_4(\psi^1, \eta, S^{(+)})=+$; $Z_4(\psi^{2,3}, S^{(0)})=-$; $Z_4(\rho, \chi) = i$. The interaction, $\eta \eta S^{(+c)} S^{(+c)}$, which is Z_4 (and L)-symmetric for $S^{(+11)}$, does not disturb $v_{\nu}^{(+)}=0$ and breaks the L' conservation but preserves a discrete symmetry based on $\exp(iL'\pi/2) (\equiv Z_{L'})$ since L' is broken by $|L'| = 4$. Then, Z_4 and $Z_{L'}$ symmetries collaborate to ensure the vacuum alignment and all possible L -violating interactions that can generate $v_{\nu}^{(+)}$ turn out to be forbidden.

The Yukawa interactions consist of $f_{1i} \overline{(\psi^1)^C} \psi^i S^{(0)}$ and $(1/2) f_{ij} \overline{(\psi^i)^C} \psi^j S^{(+)}$ for $i, j = 2, 3$ (without $\psi \psi \eta$ forbidden by Z_4) and Higgs potentials are given by trivial self-Hermitian terms and by non self-Hermitian terms composed of $\mu_0 \rho S^{(0)} \chi$ and $\lambda_1 \eta \eta S^{(0)C} S^{(0)C}$ generating $v_{\ell}^{(0)} \neq 0$ and $\lambda_2 \eta S^{(0)C} \rho^C \chi^C$ generating $v_{\nu}^{(0)} \neq 0$, both of which conserve L' , and of $\lambda_b \eta \eta S^{(+C)} S^{(+C)}$ generating $v_{\ell}^{(+)} \neq 0$, which violates L' . These Higgs interactions support our vacuum alignment.

We obtain neutrino and charged lepton mass matrices as follows:

$$M_{\nu} = \begin{pmatrix} \delta m_{11}^{rad} & \epsilon \delta m_{12}^{\ell} & \epsilon \delta m_{13}^{\ell} \\ \epsilon \delta m_{12}^{\ell} & 0 & 0 \\ \epsilon \delta m_{13}^{\ell} & 0 & 0 \end{pmatrix}, \quad M_{\ell} = \begin{pmatrix} 0 & \delta m_{12}^{\ell} & \delta m_{13}^{\ell} \\ \delta m_{12}^{\ell} & m_{22}^{\ell} & m_{23}^{\ell} \\ \delta m_{13}^{\ell} & m_{23}^{\ell} & m_{33}^{\ell} \end{pmatrix} \quad (1)$$

where δm_{1i}^{ℓ} (m_{ij}^{ℓ}) = $f_{1i} v_{\ell}^{(0)}$ ($f_{ij} v_{\ell}^{(+)}$) ($i, j=2,3$), $\epsilon = v_{\nu}^{(0)}/v_{\ell}^{(0)}$, δm_{11}^{rad} is a radiative mass as in Fig.1: $\delta m_{11}^{rad} \sim -2\lambda_2 f_{13}^2 m_{\tau} \mu_0 v_{\chi}^2 / v_{weak}^3$ ($v_{weak} \equiv 174$ GeV). The hierarchical parameterization of m_{ij}^{ℓ} is used for $m_{\mu} \ll m_{\tau}$ and $\delta m_{1i}^{\ell} \ll m_{ij}^{\ell}$ for $m_e \ll m_{\mu, \tau}$ [7] and the bimaximal structure relies upon $\delta m_{12}^{\ell} \approx \delta m_{13}^{\ell}$.

Using the phenomenologically acceptable parameter set of ($\mu_0 = m_\chi = v_\chi$, $v_{\eta,\rho,S^{(0,+)}} = m_{\rho,S^{(0,+)}} = v_{weak}/2$) and an estimate of $v_\nu^{(0)} \sim -\lambda_2 v_\eta v_\rho / v_\chi$ [8] with $\lambda_2 \sim 10^{-7}$ for $m_\nu \sim 5 \times 10^{-3}$ eV², we finally obtain $\Delta m_\odot^2 / \Delta m_{atm}^2 \sim 2\delta m_{11}^{rad} / m_\nu \sim (10^{-2} - 10^{-1})$ for $v_\chi \sim 3.5 - 6.0$ TeV, which are consistent with the $SU(3)_L$ breaking scale. This model is, thus, relevant to yield the LMA solution.¹

We have constructed the minimal $SU(3)_L \times U(1)_N$ model with the leptons placed in (ν, ℓ, ℓ^C) . The approximate bimaximal mixings are based on the L' symmetry and the specific emphasis is laid on the fact that the similarity of our neutrino mass matrix to the charged lepton mass matrix for the (1,2) and (1,3)-entries in the ϵ terms of Eq.(1). We found sufficient power of symmetries that forbids neutrino mass terms in the (i, j) -entries ($i, j = 2, 3$) while these terms are allowed for charged leptons. The observed mass hierarchy $\Delta m_{atm}^2 \gg \Delta m_\odot^2$ is explained by the dynamical one between tree and one-loop effects and the LMA solution shows up for $v_\chi \sim$ a few TeV.

The work of M.Y. is supported by the Grant-in-Aid for Scientific Research No 12047223 from the Ministry of Education, Science, Sports and Culture, Japan.

References

- [1] Z. Maki, M. Nakagawa and S. Sakata, Prog. Theor. Phys. **28** (1962) 870.
- [2] F. Pisano and V. Pleitez, Phys. Rev. D **46** (1992) 410; P.H. Frampton, Phys. Rev. Lett. **69** (1992) 2889; D. Ng, Phys. Rev. D **49** (1994) 4805; M.Özer, Phys. Rev. D **54** (1996) 1143. See also, M. Singer, J.W.F. Valle and J. Schechter, Phys. Rev. D **22** (1980) 738.
- [3] T. Kitabayashi and M. Yasuè, hep-ph/0103265 (Mar.,2001) to be published in Nucl. Phys. B (2001). For other variations, see, T. Kitabayashi and M. Yasuè, Phys. Rev. D **63** (2001) 095002 and 095006; Phys. Lett. B **503** (2001) 85; T. Kitabayashi, hep-ph/0103195 to be published in Phys. Rev. D **64** (2001).
- [4] See for example, K. Nishikawa, a talk at the international Workshop NuFACT'01, Tsukuba Japan (May 2001); Y. Fukuda, a talk at NuFACT'01.
- [5] S.T. Petcov, Phys. Lett. **110B** (1982) 245; C.N. Leung and S. T. Petcov, Phys. Lett. **125B** (1983) 461; A. Zee, Nucl. Phys. **264B** (1986) 99; R. Barbieri, L.J. Hall, D. Smith, N.J. Weiner and A. Strumia, JHEP **12** (1998) 017.
- [6] A. Zee, Phys. Lett. **93B** (1980) 389; Phys. Lett. **161B** (1985) 141; L. Wolfenstein, Nucl. Phys. B **175** (1980) 93.

¹ The ratio $\Delta m_\odot^2 / \Delta m_{atm}^2$ is also induced by the diagonalization effects of M_ℓ estimated to be $\lesssim (m_e/m_\mu)^{3/2} \approx 3.4 \times 10^{-4}$, which can be safely neglected [3].

- [7] See for example, H. Fritzsch, Phys. Lett. **70B** (1977) 436; Phys. Lett. **73B** (1978) 317. For our analysis in other cases, see Ref.[3] for the details.
- [8] M.B. Tully and G.C. Joshi, Phys. Rev. D **64** (2001) 011301(R).